

# sul-fa-meth-OX-a-zole, trye-METH-oh-prim Brand Name: Septra, Sulfatrim, Bactrim

Sulfamethoxazole-trimethoprim, also known as Bactrim, Septra, or TMP-SMX, belongs to the classes of medicines called antibacterials and antiprotozoals. Antibacterials kill bacteria or stop bacteria from multiplying. Antiprotozoals help the body overcome infections caused by protozoa, microscopic organisms that can cause disease. When bacterial and protozoal infections occur in patients with weakened immune systems, including people with HIV or AIDS, they tend to be more serious and are called opportunistic infections (OIs).

# **HIV/AIDS-Related Uses**

Sulfamethoxazole-trimethoprim was approved by the FDA on June 23, 1981, for use in the treatment and prophylaxis of Pneumocystis carinii pneumonia (PCP), a common OI in people with HIV.

## **Dosage Information**

Sulfamethoxazole-trimethoprim comes in oral suspension, tablet, and injectable forms. The oral suspension and tablet forms of sulfamethoxazole-trimethoprim should be taken with a full glass of water, and the oral suspension should be shaken well before use.

#### **Contraindications**

Individuals should tell a doctor about any medical problems before taking this medicine.

#### **Possible Side Effects**

Along with its desired effects, sulfamethoxazole-trimethoprim can cause some unwanted effects. Serious side effects of this medicine include itching; skin rash; aching of joints and muscles; difficulty in swallowing; pale skin; redness; blistering, peeling, or loosening of skin; sore throat and fever; unusual bleeding or bruising; unusual tiredness or weakness; yellow eyes or skin; or increased sensitivity of skin to sunlight. Individuals should tell a doctor if they have symptoms of any of these side effects.

Other side effects may not be serious and may lessen or disappear with continued use of the medicine. Less serious side effects of this medicine include diarrhea, dizziness, headache, loss of appetite, mouth sores or swelling of the tongue, nausea or vomiting, and tiredness. Individuals should tell a doctor if these side effects continue or are bothersome.

### **Drug and Food Interactions**

A doctor should be notified of any other medications being taken, including prescription, nonprescription (over-the-counter), or herbal medications.

#### **Clinical Trials**

For information on clinical trials that involve Sulfamethoxazole-Trimethoprim, visit the ClinicalTrials.gov web site at http://www.clinicaltrials.gov. In the Search box, enter: Sulfamethoxazole-Trimethoprim AND HIV Infections.

## **Manufacturer Information**

Sulfamethoxazole-Trimethoprim King Pharmaceuticals, Inc. 501 Fifth Street Bristol, TN 37620 (888) 840-5370

#### Septra

King Pharmaceuticals, Inc. 501 Fifth Street Bristol, TN 37620 (888) 840-5370

Sulfatrim Alpharma, Inc. One Executive Drive Fort Lee, NJ 07024 (201) 947-7774

# Bactrim

Women First Healthcare 5355 Mira Sorrento Place - Suite 700 San Diego, CA 92121 (858) 509-1171



# **For More Information**

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday Friday, 12:00 p.m. (Noon) 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live\_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET